

PERIOD-INDEX IN TOP COHOMOLOGY OVER SEMIGLOBAL FIELDS

SARAH DIJOLS, RAMAN PARIMALA, SUJATHA RAMDORAI, AND CHARLOTTE URE

ABSTRACT. We prove a common slot lemma for symbols in top cohomology classes over semiglobal fields. Furthermore, we prove that period and index agree for general top cohomology classes over such fields. We discuss applications to quadratic forms and related open problems.

1. INTRODUCTION

Let F be a field and fix a separable closure \overline{F} of F . The Galois group $\text{Gal}(\overline{F}/F)$ will be denoted by G_F . Let μ_m be the group of m -th roots of unity in \overline{F} , considered as a module over G_F , and denote by $\mu_m^{\otimes i}$ its i -fold tensor product. The Galois cohomology groups $H^i(G_F, \mu_m^{\otimes i})$ will be denoted $H^i(F, \mu_m^{\otimes i})$ throughout. For any field extension E of F , the profinite group G_E is a subgroup of G_F and so there are natural restriction maps

$$\text{res} : H^i(F, \mu_m^{\otimes i}) \rightarrow H^i(E, \mu_m^{\otimes i}). \quad (1.1)$$

that take a cohomology class ξ to its restriction ξ_E . If $\xi_E = 0$, we call E a splitting field of ξ .

We call an element $\xi \in H^i(F, \mu_m^{\otimes i})$ a *symbol* if it is in the image of the i -fold cup product map

$$H^1(F, \mu_m) \times \cdots \times H^1(F, \mu_m) \rightarrow H^i(F, \mu_m^{\otimes i}).$$

If $\mu_m \subset F$, then $\mu_m^{\otimes i} \cong \mu_m$ as G_F -modules, so that there is a notion of symbols in $H^i(F, \mu_m)$. In the case that $i = m = 2$, a symbol in $H^2(F, \mu_2^{\otimes 2})$ determines a quaternion algebra over F and vice versa. Tate's common slot lemma states that if F is a number field, then any finite set of symbols in $H^2(F, \mu_m^{\otimes 2})$ can be split by a single extension F' of degree dividing m [Tat76, Theorem 4.4 and the remarks following the theorem]. In particular, any finite set of quaternion algebras over a number field may be split by a single quadratic extension of F . A semiglobal field F is a function field

Date: May 26, 2024.

2020 Mathematics Subject Classification. Primary 12G05; Secondary 11E04, 11E81, 13F25, 14F22, 14H25.

Key words and phrases. Galois cohomology, symbol length, patching, semiglobal fields.

in one variable over a complete discretely valued field K . We prove an analogue of Tate's common slot lemma in the setting of semiglobal fields in Theorem 4.3.

Understanding the splitting fields of general cohomology classes, as in Tate's common slot lemma, has been an interesting problem in the study of Galois cohomology. This motivates the notion of the index of $\xi \in H^i(F, \mu_m)$ as the greatest common divisor of the degrees of field extensions E of F so that ξ splits over E . For an element ξ in $H^i(F, \mu_m)$, we can also define the period of ξ to be its order, which is denoted by $\text{per}(\xi)$. It is known that the period divides the index and that they have the same prime factors ([Pie82], Proposition 14.4(b)(ii)). The period-index question asks for the smallest number r so that

$$\text{ind}(\xi) \mid \text{per}(\xi)^r$$

for all $\xi \in H^i(F, \mu_m)$. There has been much progress toward answering this problem over various fields. In the case that F is a local or global field, period and index agree in the Brauer group ($i = 2$) by the Albert–Brauer–Hasse–Noether Theorem [BNH32]. Recent important results have been obtained in the case of degree 3 cohomology, where [PS98] showed that period and index coincide in $H^3(F, \mu_m)$ if F is the function field of a p -adic curve or the function field of a surface over a finite field. Furthermore, this coincidence has been settled for function fields of curves over imaginary number fields in [Sur20]. In [HHK23], the authors obtain bounds on the index in terms of the period for F a semiglobal field.

Throughout this paper, F is a semiglobal field, i.e. a function field in one variable over a complete discretely valued field K with residue field k . Let m be a number coprime to the characteristic of k , and n be the largest number such that $H^n(k, \mu_m)$ does not vanish. We investigate the cohomology groups $H^{n+2}(F, \mu_m)$. Since $H^i(F, \mu_m)$ vanishes for any $i > n + 2$, we call $H^{n+2}(F, \mu_m)$ the *top cohomology*. For classes $\xi \in H^{n+2}(F, \mu_m)$, Gosavi proved in [Gos22, Corollary 3.3] that $\text{ind}(\xi) \mid \text{per}(\xi)^2$ for m an odd prime and $\text{ind}(\xi) \mid \text{per}(\xi)^3$ for $m = 2$. In Theorem 4.8, we prove that period and index agree in top cohomology for semiglobal fields for any m coprime to the characteristic of k . In fact, for a class $\xi \in H^{n+2}(F, \mu_m)$, we construct a field extension L of F of degree dividing m that is a splitting field of ξ .

This paper is organized as follows. In Section 2, we give necessary background on cohomological dimension, the period-index problem, semiglobal fields, and the local-global principle. Let K be a complete discretely valued field with residue field k of characteristic coprime to m and suppose that k has cohomological dimension n . In Section 3, we explore the top cohomology groups $H^{n+1}(K, \mu_m)$ of the complete discretely valued field K . In particular, we prove Lemma 3.1, which provides a decomposition of elements in $H^i(K, \mu_m)$ that is crucial to our proofs. Using this decomposition, we show that period and index agree in the top cohomology $H^{n+1}(K, \mu_m)$

groups in Lemma 3.4. In Section 4, we state and prove our two main theorems. First, we prove an analogue of Tate's common slot lemma over semiglobal fields in Theorem 4.3. We show that for any finite set \mathcal{S} of symbols in $H^{n+2}(F, \mu_m^{\otimes n+2})$, there is a field extension L of F of degree dividing m so that all $s \in \mathcal{S}$ are split over L . The proof relies on the local-global principle discussed in Theorem 2.11. More precisely, we show that for any discrete valuation ω on L with restriction ν to F , the restriction s_{F_ν} splits over s_{L_ω} for all $s \in \mathcal{S}$. Finally, we show in Theorem 4.8 that period and index agree in top cohomology of $H^{n+2}(F, \mu_m)$. We conclude the paper by discussing some open questions on top cohomology and connections of period-index questions with the u -invariant of quadratic forms in Section 5.

Acknowledgement. First, we thank Suresh Venapally for bringing to our attention the questions on top cohomology and for continued support during the project. We also thank Danny Krashen for helpful comments and suggestions. We thank the referee for their careful reading and suggestions.

This project was started during the Women in Numbers (WIN) workshop at the Banff International Research Station in 2023. We thank the workshop organizers Shabnam Akhtari, Alina Bucur, Jennifer Park, Renate Scheidler, and the staff at BIRS for their support. We also thank the following for supporting the workshop, childcare, and travel: NSF DMS-2012061, the Number Theory Foundation, the Clay Mathematics Institute, the Journal of Number Theory, the Pacific Institute for Mathematical Sciences, and the AMS-Simons Travel Grant program. Sujatha Ramdorai is supported by the NSERC Discovery Grant 2019-03987.

2. PRELIMINARIES

2.1. Cohomological dimension. For a profinite group G and a G -module A , we denote by $H^i(G, A)$ the i -th profinite cohomology group. If G_F is the absolute Galois group of a field F , we also call A an F -module and write $H^i(F, A) = H^i(G_F, A)$ (see for instance [Ser73, Chapter I §2.2 and Chapter II]). The cohomological dimension is an important invariant associated with a field (for more details, we refer the reader to [Ser73, Chapter I §3 and Chapter II §4]).

Definition 2.1. *For a prime ℓ , the ℓ -cohomological dimension of F , $\text{cd}_\ell(F)$, is the smallest integer R for which the ℓ -primary component of $H^{i+R}(F, A)$ is zero for any integer $i \geq 1$ and for every torsion F -module A . If no such R exists, we say that $\text{cd}_\ell(F) = \infty$. Analogously, the cohomological dimension $\text{cd}(F)$ of F is the smallest integer R so that $H^{i+R}(F, A) = 0$ for any $i \geq 1$ and for any torsion F -module A . If no such R exists, we say that $\text{cd}(F) = \infty$.*

In this paper, we are only interested in the groups $H^i(F, \mu_m)$ for $m \in \mathbb{Z}$, $m \geq 2$, $(m, \text{char}(F)) = 1$. We remark that if the characteristic of F is different from ℓ , for ℓ

prime, then by [Ser73, Proposition 4, Chapter II, §2.3] the following are equivalent for any $n \geq 1$:

- (a) $\text{cd}_\ell(F) \leq n$;
- (b) For any finite separable extension L of F of degree prime to ℓ , $H^{n+1}(L, \mu_\ell) = 0$

An algebraically closed field has cohomological dimension 0. Finite fields and the Laurent series field $k((t))$ with k algebraically closed of characteristic 0 have cohomological dimension 1. More generally, the cohomological dimension is invariant under finite field extensions (see Lemma 2.2 below) and increases as the transcendental degree increases (see Lemma 2.3 below).

Lemma 2.2 ([Ser73, Proposition 10, Chapter II, §4.1]). *Let L be an algebraic extension of the field F and ℓ a prime number. Assume $[L : F] < \infty$ and $\text{cd}_\ell(F) < \infty$, then $\text{cd}_\ell(L) = \text{cd}_\ell(F)$. In particular, if $[L : F] < \infty$ and $\text{cd}(F) < \infty$, then $\text{cd}(L) = \text{cd}(F)$.*

Lemma 2.3 ([Ser73, Proposition 11, Chapter II, §4.2]). *Let L be an extension of K , then*

$$\text{cd}_\ell(L) \leq \text{cd}_\ell(K) + \text{trdeg}_K(L)$$

for any prime ℓ . There is equality if L is finitely generated over K , $\text{cd}_\ell(K) < \infty$, and ℓ is coprime to the characteristic of K .

Let K be a complete discretely valued field with residue field k and suppose that $\text{char}(k) \neq \ell$. For $n \geq 1$, we consider the following exact sequence in Galois cohomology

$$0 \rightarrow H^{n+1}(k, \mu_\ell) \rightarrow H^{n+1}(K, \mu_\ell) \rightarrow H^n(k, \mathbb{Z}/\ell\mathbb{Z}) \rightarrow 0. \quad (2.1)$$

This sequence is obtained combining the surjective homomorphism between $H^n(k, \mu_\ell)$ and $H^n(k, \mathbb{Z}/\ell\mathbb{Z})$ given in 4.3 [Ser97] and residue maps (see 6.8 [GS17]). Using this sequence, we deduce the following relationship between the cohomological dimension of K and k , respectively.

Lemma 2.4. *Let K be a complete discretely valued field with residue field k and suppose that $\text{char}(k) \neq \ell$. If $\text{cd}_\ell(k) \leq n$, then $\text{cd}_\ell(K) \leq n + 1$.*

Proof. Let L be a finite separable extension of K of degree coprime to ℓ with residue field l , containing k . As $\text{cd}(k) \leq n$ and l is a finite extension of k of degree coprime to ℓ , we deduce from Lemma 2.2 that $\text{cd}_\ell(l) = \text{cd}_\ell(k) \leq n$. In particular, we see that $H^{n+1}(l, \mathbb{Z}/\ell\mathbb{Z}) = 0$. The sequence in (2.1) implies that $H^{n+2}(L, \mu_\ell) = 0$. As this holds for any finite extension L of K , we conclude that $\text{cd}(K) \leq n + 1$. \square

Lemma 2.5. *Let K be a complete discretely valued field with residue field k and suppose that $\text{char}(k) \neq \ell$. If $\text{cd}_\ell(k) = n$, then $\text{cd}_\ell(K) = n + 1$.*

Proof. By Lemma 2.4, it remains to show that $\text{cd}_\ell(K) \geq n + 1$, or equivalently, that there is some finite field extension L of K of degree coprime to ℓ so that $H^{n+1}(L, \mu_\ell) \neq 0$.

By assumption $\text{cd}_\ell(k) = n$ and so there exists a finite separable field extension l of k of degree coprime to ℓ so that $H^n(l, \mu_\ell) \neq 0$. Let ρ be a primitive ℓ -th root of unity in the algebraic closure of l . We remark that $l[\rho]$ is a separable extension of l of degree coprime to ℓ . Replacing l by $l[\rho]$, we may assume without loss of generality, that ρ is contained in l . Therefore

$$H^n(l, \mathbb{Z}/\ell\mathbb{Z}) = H^n(l, \mu_\ell) \neq 0. \quad (2.2)$$

By the primitive element theorem, there is some $\bar{\alpha} \in l$ so that $l = k(\bar{\alpha})$. Denote by $\bar{f}(x) \in k[x]$ the minimal polynomial of $\bar{\alpha}$ over k . Let $f(x) \in K[x]$ be a monic lift of $\bar{f}(x)$ and consider $L = K(\alpha) = K[x]/(f(x))$. Then L/K is a separable extension of degree coprime to ℓ that lifts l/k . The short exact sequence in (2.1) for L and l together with equation (2.2) implies that $H^{n+1}(L, \mu_\ell)$ does not vanish. \square

Let $F = K(C)$ be the function field of an algebraic curve over a complete discretely valued field K with residue field k . Then F is a semiglobal field in the sense discussed below in Section 2.3. If ℓ is a prime number which is coprime to the characteristic of the residue field k , then it is also coprime to the characteristic of K and F . In particular, we may apply Lemmas 2.3 and 2.5 to F and K in this setting to deduce that if $\text{cd}_\ell(k) = n$, then $\text{cd}_\ell(K) = n + 1$ and $\text{cd}_\ell(F) = n + 2$. Furthermore, if $\text{cd}(k) = n$, then $\text{cd}_\ell(K) \leq n + 1$ and $\text{cd}_\ell(F) \leq n + 2$ for any prime ℓ coprime to the residue characteristic.

For a field F with $\text{cd}(F) = n$ and A a torsion G -module, the *top cohomology with coefficients in A* is the cohomology group $H^n(F, A)$. Throughout this paper, A will often be the Galois module arising from roots of unity.

2.2. Period-index problem and Voevodsky's Theorem. Let F be a field and let m be any number coprime to the characteristic of F . Fix a separable closure \bar{F} of F and denote by G_F its absolute Galois group. Consider the Kummer exact sequence of G_F -modules

$$1 \longrightarrow \mu_m \longrightarrow \bar{F}^\times \xrightarrow{a \mapsto a^m} \bar{F}^\times \longrightarrow 1.$$

The induced long exact sequence of cohomology in conjunction with Hilbert 90 gives an isomorphism

$$H^1(F, \mu_m) \xrightarrow{\sim} F^\times / (F^\times)^m. \quad (2.3)$$

There is an i -fold cup-product from the product of i copies of $H^1(F, \mu_m)$ to $H^i(F, \mu_m)$ for any $i > 1$

$$\cup : H^1(F, \mu_m) \times \cdots \times H^1(F, \mu_m) \longrightarrow H^i(F, \mu_m). \quad (2.4)$$

For elements $a_1, \dots, a_i \in F^\times / (F^\times)^m$, we denote by $(a_1, \dots, a_i) \in H^i(F, \mu_m)$ the image under the composition of the isomorphism in (2.3) and the cup product in (2.4). Such an element in $H^i(F, \mu_m)$ is called a *symbol*. Abusing notation, we will consider $a_1, \dots, a_i \in F^\times$ throughout. The cup-product map in (2.4) is alternating and we have the following relations for any $a, b \in F^\times$ and $n \in \mathbb{Z}$

$$\begin{aligned} -(a, b) &= (b, a) \\ n(a, b) &= (a, b^n) = (a^n, b) \\ (a, a) &= (-1, a) \end{aligned} \quad (2.5)$$

(see e.g. [GS17, Chapter 3.4 and Chapter 4.6]).

Let $K_i^M(F)$ denote the i -th Milnor K -group of F (see H. Bass's Algebraic K-theory). Let ℓ be a prime not equal to $\text{char}(F)$. Suppose that F contains the ℓ -th roots of unity μ_ℓ . By a celebrated result of Voevodsky [Voe03], the norm-residue map determines an isomorphism between Milnor's K -groups and cohomology

$$K_i^M(F) \otimes \mathbb{Z}/\ell\mathbb{Z} \xrightarrow{\sim} H^i(F, \mu_\ell).$$

In particular, any element ξ in $H^i(F, \mu_\ell)$ may be written as a sum of symbols

$$\xi = \sum_{j=1}^r (a_{j,1}, \dots, a_{j,i}),$$

with $a_{j,1}, \dots, a_{j,i} \in F^\times$ for $1 \leq j \leq r$. Related to this presentation, the *symbol length* of the i -th cohomology group $H^i(F, \mu_\ell)$, denoted $\text{symb}_i^\ell(F)$, is defined as the minimal number r such that any element $\xi \in H^i(F, \mu_\ell)$ is expressible as a sum of at most r symbols. Investigating the symbol lengths $\text{symb}_i^\ell(F)$ for fields F is of interest. For more background and open problems related to the symbol lengths of fields, we refer the interested reader to [ABGV11]. We note that by definition, the symbol length $\text{symb}_1^\ell(F) = 1$ for any field F . For $n = 2$, the second cohomology group $H^2(F, \mu_\ell)$ is isomorphic to the ℓ -torsion of the Brauer group $\text{Br}(F)$. For a p -adic local field F , the Brauer group $\text{Br}(F) \cong \mathbb{Q}/\mathbb{Z}$ [Ser97, Chapter XII] and if $\mu_\ell \subset F$, then $\text{symb}_2^\ell(F) = 1$. Using this together with the Albert–Brauer–Hasse–Noether Theorem [BNH32], we deduce that $\text{symb}_2^\ell(F) = 1$ for any totally imaginary global field F . In either of the two previous cases $\text{cd}(F) = 2$ and so in summary

$$\text{symb}_n^\ell(F) = \begin{cases} 1 & n = 1, 2 \\ 0 & n > 2 \end{cases}$$

for a local or a totally imaginary global field F containing μ_ℓ . Now, let F be the function field of a p -adic curve, and let ℓ be a prime which is coprime to p , then the symbol length $\text{symb}_2^\ell(F)$ is 2 [Sal97, Sal98] and $\text{symb}_3^\ell(F) = 1$ [PS10].

Remark 2.6. *We recall that a cohomology class $\xi \in H^i(F, A)$ for a field F and a Galois module A has splitting field L if the restriction ξ_L as defined in (1.1) is trivial. For example, any symbol (a, b) in $H^2(F, \mu_m)$ is split by the degree m extension $L = F(\sqrt[m]{a})$ since*

$$0 = m \left(\sqrt[m]{a}, b \right)_L \stackrel{(2.5)}{=} \left(\left(\sqrt[m]{a} \right)^m, b \right)_L = (a, b)_L,$$

where the first equality holds as $H^n(L, \mu_m)$ is m -torsion. More generally, a symbol $(a_1, \dots, a_i) \in H^i(F, \mu_m)$ splits over the degree m extension $F(\sqrt[m]{a_j})$ for any $1 \leq j \leq i$.

The index of a cohomology class gives a measure for splitting.

Definition 2.7. *Let F be a field and $\xi \in H^i(F, A)$. The index of ξ is*

$$\text{ind}(\xi) = \text{gcd} \{ [L : F] : \xi_L \text{ splits} \}.$$

The period-index question discussed in the introduction explores the relationship between symbol lengths and the degrees of splitting field extensions for symbols. In Section 5.2 we will explain the connection between symbol lengths and the u -invariant of a field.

2.3. Semiglobal fields and a local-global principle. One of the main tools we build on in this paper is patching in semiglobal fields. We follow [HHK14] in the following definitions.

Definition 2.8 (semiglobal fields). *Let K be a complete discretely valued field. A semiglobal field over K is a one-variable function field F over K .*

Example 2.9. *Examples of semiglobal fields are $F = \mathbb{Q}_p(x)$, $F = k((t))(x)$ for any field k , and any finite extension of these.*

Let F be a semiglobal field, that is F is the function field of a smooth projective curve C over K . Let \mathcal{O}_K be the corresponding valuation ring, $\pi \in \mathcal{O}_K$ a (fixed) uniformizing parameter and k the residue field. A *regular proper model* of C is a proper, flat, projective \mathcal{O}_K -scheme \mathfrak{X} with function field F that is regular as a scheme. We denote its reduced closed fiber by \mathfrak{X}_k . By [Lip78, Main Theorem] such a regular proper model exists and by [Lip75, page 193], we may assume that the reduced closed fiber \mathfrak{X}_k is a union of regular curves with normal crossings. We call

such a model a *normal crossings model* of F .

$$\begin{array}{ccccc} C & \longrightarrow & \mathfrak{X} & \longleftarrow & \mathfrak{X}_k \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{Spec}(K) & \longrightarrow & \mathrm{Spec}(\mathcal{O}_K) & \longleftarrow & \mathrm{Spec}(k) \end{array}$$

For each prime divisor D , we can look at the discretely valued ring $\mathcal{O}_{D,\mathfrak{X}}$ and get a corresponding discrete valuation v_D on F . Such valuations are known as *divisorial*.

Definition 2.10. *The divisorial set of places of F corresponding to the model \mathfrak{X} is the set*

$$\Omega_{F,\mathfrak{X}} = \{v_D \mid D \text{ prime divisor on } \mathfrak{X}\}.$$

We call v_D with D as above a *divisorial valuation* on F . The set of all divisorial valuations v_D , on all regular models \mathfrak{X} of F , will be denoted by Ω_F .

A new approach to local-global principles for homogeneous varieties over function fields of curves defined over complete discretely valued fields was introduced by Harbater, Hartmann, and Krashen in [HHK14] via patching. Further, Parimala and Suresh, building on the results of Hartmann, Harbater, and Krashen, proved the following Theorem that will fundamentally be used later in our paper:

Theorem 2.11 (Theorem 4.3, [HKP21]). *Let F be a semiglobal field, ℓ be a prime number, Ω_F the set of divisorial places of F , and let r be any integer. Suppose that $\mathrm{char}(k) \neq \ell$, and let $i \geq 2$. Then*

$$\mathrm{III}_{\Omega_F}^i := \mathrm{Ker} \left(H^i(F, \mu_\ell^{\otimes r}) \rightarrow \prod_{\omega \in \Omega_F} H^i(F_\omega, \mu_\ell^{\otimes r}) \right) = 0.$$

3. COMPLETE DISCRETELY VALUED FIELD OF COHOMOLOGICAL DIMENSION $n + 1$

Let K be a complete discretely valued field, \mathcal{O}_K its ring of integers, π its local uniformizer, and k its residue field. Throughout this section, we suppose that ℓ is a prime which is coprime to the residue characteristic and $\mathrm{cd}_\ell(k) = n$. By Lemma 2.5, we see that $\mathrm{cd}_\ell(K) = n + 1$. As a result, the expression *top cohomology* of K will refer to $H^{n+1}(K, \mu_\ell)$. The aim of this section is to prove that period and index agree in top cohomology in this setting.

We first assume that $\mu_\ell \subset k$, which implies that $\mu_\ell \subset K$ by Hensel's lemma, using that ℓ is coprime to the characteristic of k . We recall that there is an exact sequence on Galois cohomology

$$0 \longrightarrow H_{ct}^{n+1}(\mathcal{O}_K, \mu_\ell) \xrightarrow{r} H^{n+1}(K, \mu_\ell) \xrightarrow{\delta} H^n(k, \mu_\ell), \quad (3.1)$$

where r is induced by restriction and δ is a residue map (see e.g. [GS17, Chapter 6.8]). An element ξ in $H^{n+1}(K, \mu_\ell)$ is called *unramified* if it lies in the kernel of δ , or equivalently in the image of r . Otherwise, we say that ξ is *ramified*.

We show that any cohomology class splits into a *sum of ramified and unramified parts*. We recall from Section 2.2 that there is a cup-product map from $\otimes^m H^1(K, \mu_\ell)$ to $H^m(K, \mu_\ell)$ and that $H^1(K, \mu_\ell) \cong K^\times / (K^\times)^\ell$.

Lemma 3.1. *Let K be a discrete valued field, \mathcal{O}_K its ring of integers, and k its residue field. Suppose that $\mu_\ell \subset k$ and denote by π the local uniformizer in \mathcal{O}_K . Any element $\xi \in H^m(K, \mu_\ell)$ decomposes as*

$$\xi = \xi_1 + (\xi_2, \pi)$$

with $\xi_1 \in H^m(K, \mu_\ell)$ and $\xi_2 \in H^{m-1}(K, \mu_\ell)$ unramified for any $m > 0$. In the above equation (ξ_2, π) denotes the cup product of ξ_2 with the symbol $(\pi) \in H^1(K, \mu_\ell)$.

Proof. Let $\xi \in H^m(K, \mu_\ell)$ and suppose first that ξ is a symbol, that is $\xi = (a_1, \dots, a_m)$ with $a_i \in K^\times$. Write $a_i = u_i \pi^{j_i}$ with u_i a unit in \mathcal{O}_K and $0 \leq j_i \leq \ell - 1$. Using the relations in (2.5), we see that

$$\begin{aligned} \xi &= (a_1, \dots, a_m) \\ &= (u_1 \pi^{j_1}, \dots, u_m \pi^{j_m}) \\ &= (u_1, u_2 \pi^{j_2}, \dots, u_m \pi^{j_m}) + (\pi^{j_1}, u_2 \pi^{j_2}, \dots, u_m \pi^{j_m}) \\ &= (u_1, u_2, u_3 \pi^{j_3}, \dots, u_m \pi^{j_m}) + (u_1, \pi^{j_2}, u_3 \pi^{j_3}, \dots, u_m \pi^{j_m}) \\ &\quad + (\pi^{j_1}, u_2, u_3 \pi^{j_3}, \dots, u_m \pi^{j_m}) + (\pi^{j_1}, \pi^{j_2}, u_3 \pi^{j_3}, \dots, u_m \pi^{j_m}) \\ &= (u_1, u_2, u_3 \pi^{j_3}, \dots, u_m^{j_m}) + \left(\frac{u_1^{j_2}}{u_2^{j_1}}, \pi, u_3 \pi^{j_3}, \dots, u_m \pi^{j_m} \right) \\ &\quad + ((-1)^{j_1+j_2}, \pi, u_3 \pi^{j_3}, \dots, u_m \pi^{j_m}) \\ &\quad \vdots \\ &= (u_1, u_2, \dots, u_m) + \sum_t (v_1^t, v_2^t, \dots, v_{m-1}^t, \pi) \end{aligned}$$

for some units v_s^t in \mathcal{O}_K . Since $\mu_\ell \subset k$, any cohomology class $\xi \in H^{n+1}(K, \mu_\ell)$ may be decomposed as a sum of symbols $\xi = \sum_i (a_{i,1}, \dots, a_{i,m})$ by Voevodsky's Theorem (see Section 2.2). We decompose each summand of ξ as above to write

$$\xi = \sum_i (a_{i,1}, \dots, a_{i,m}) = \sum_i (u_{i,1}, \dots, u_{i,m}) + \sum_{i,t} (v_{i,1}^t, \dots, v_{i,m-1}^t, \pi) = \xi_1 + (\xi_2, \pi),$$

which finishes the proof. \square

Using the description from the previous Lemma, we are now able to explicitly determine a splitting field of $\xi \in H^{n+1}(K, \mu_\ell)$. We recall from Remark 2.6, that a symbol $(a_1, \dots, a_n) \in H^n(K, \mu_\ell)$ may be split by the extension $L = K(\sqrt[\ell]{a_i})$ for any $1 \leq i \leq n$.

Lemma 3.2. *Let K be a complete discretely valued field, π its local uniformizer, and k its residue field. Suppose that $\text{cd}_\ell(k) = n$. Assume additionally that $\mu_\ell \subset k$. Then any element $\xi \in H^{n+1}(K, \mu_\ell)$ is split by the degree ℓ field extension $K(\sqrt[\ell]{\pi})$ of K . In particular, it has index bounded above by ℓ .*

Proof. We note that as $\text{cd}_\ell(k) = n$, by Lemma 2.5, we have $\text{cd}_\ell(K) = n + 1$. Let $\xi \in H^{n+1}(K, \mu_\ell)$. By Lemma 3.1, ξ can be written as $\xi = \xi_1 + (\xi_2, \pi)$ for some unramified $\xi_1 \in H_{\text{et}}^{n+1}(\mathcal{O}_K, \mu_\ell)$ and $\xi_2 \in H_{\text{et}}^n(\mathcal{O}_K, \mu_\ell)$. We note that if ξ is unramified, then $\xi = \xi_1$. Since \mathcal{O}_K is Henselian, we remark that $H_{\text{et}}^m(\mathcal{O}_K, \mu_\ell) \cong H^m(k, \mu_\ell)$ for any m . Together with the fact that $\text{cd}_\ell(k) = n$, we deduce that $\xi_1 = 0$. In particular, if ξ is unramified, then $\xi = \xi_1 = 0$ is already split over K so that $\text{ind}(\xi) = 0 \leq \ell$. In general, we write any $\xi = (\xi_2, \pi)$ with ξ_2 as before. This cohomology class is split by the degree ℓ field extension $K(\sqrt[\ell]{\pi})$ of K so that $\text{ind}(\xi)$ is again bounded above by ℓ . \square

The following lemma extends Lemma 3.2 to the case $\mu_\ell \not\subset k$.

Lemma 3.3. *Let K be a complete discretely valued field, π its local uniformizer, and k its residue field. Suppose that $\text{cd}_\ell(k) = n$ and assume that the characteristic of k is coprime to ℓ , but possibly $\mu_\ell \not\subset k$. Then any element $\xi \in H^{n+1}(K, \mu_\ell)$ is split by the degree ℓ field extension $K(\sqrt[\ell]{\pi})$ of K . In particular, it has index bounded above by ℓ .*

Proof. We note that as $\text{cd}_\ell(k) = n$, by Lemma 2.5, we have $\text{cd}_\ell(K) = n + 1$. Since K does not contain μ_ℓ , let us first consider the degree $\ell - 1$ extension $E = K(\rho)$ where ρ is a generator of μ_ℓ . We remark that the characteristic of k is coprime to ℓ , so that the cyclotomic extension E over K is unramified and thus π is a local uniformizer for E as well [Cas86, Equation (5.5) in the proof of Lemma 5.4]. By Lemma 3.2, after restricting to E , ξ is split by the extension $E(\sqrt[\ell]{\pi})$. Now consider the following diagram of field extensions

$$\begin{array}{ccc}
 & E(\sqrt[\ell]{\pi}) & \\
 \ell-1 \swarrow & & \searrow \ell \\
 K(\sqrt[\ell]{\pi}) & & E \\
 \searrow \ell & & \swarrow \ell-1 \\
 & K &
 \end{array}$$

The composition

$$H^{n+1}(K(\sqrt[\ell]{\pi}), \mu_\ell) \xrightarrow{\text{res}} H^{n+1}(E(\sqrt[\ell]{\pi}), \mu_\ell) \xrightarrow{\text{cor}} H^{n+1}(K(\sqrt[\ell]{\pi}), \mu_\ell)$$

is multiplication by $[E(\sqrt[\ell]{\pi}) : K(\sqrt[\ell]{\pi})] = \ell - 1$ [NSW08, Corollary 1.5.7]. Thus

$$(\ell - 1)\xi_{K(\sqrt[\ell]{\pi})} = \text{cor} \circ \text{res}(\xi_{K(\sqrt[\ell]{\pi})}) = \text{cor}(0) = 0.$$

As ℓ and $\ell - 1$ are coprime, we deduce that $\xi_{K(\sqrt[\ell]{\pi})} = 0$ as well and so $K(\sqrt[\ell]{\pi})$ is a splitting field for ξ . \square

Finally, using induction on the number of prime factors, we prove the general period-index result for complete discretely valued fields.

Lemma 3.4. *Let K be a complete discretely valued field and k its residue field. Suppose that the characteristic of k is coprime to an integer $m > 1$. Suppose that $\text{cd}(k) = n$. Then any element $\xi \in H^{n+1}(K, \mu_m)$ has index bounded above by m .*

Since m is coprime to the residue characteristic and $\text{cd}_\ell(k) \leq \text{cd}(k) = n$ for any prime divisor ℓ of m , by Lemma 2.2 and Lemma 2.5, we remark that $\text{cd}_\ell(K) \leq n + 1$ and so $H^r(K, \mu_m) = 0$ for any $r > n + 1$. This motivates the term *top cohomology* for $H^{n+1}(K, \mu_m)$ in this setting.

Proof. We use induction on the number of prime factors of m . For $m = \ell$ prime, the result follows from Lemma 3.3. If $m = s\ell$ for some prime ℓ , consider the short exact sequence of K -modules

$$0 \longrightarrow \mu_\ell \longrightarrow \mu_m \xrightarrow{\ell} \mu_s \longrightarrow 0. \quad (3.2)$$

For any $\xi \in H^{n+1}(K, \mu_m)$, let $\tilde{\xi}$ be its image under the map

$$H^{n+1}(K, \mu_m) \longrightarrow H^{n+1}(K, \mu_s)$$

arising from the long exact sequence in cohomology. By the inductive hypothesis, there is an extension L_1 of K of degree at most s such that L_1 splits $\tilde{\xi}$. By exactness of the long exact sequence of cohomology induced by (3.2), the restriction ξ_{L_1} of ξ to L_1 lifts to an element $\eta \in H^{n+1}(L_1, \mu_\ell)$.

$$\begin{array}{ccccc} H^{n+1}(L_1, \mu_\ell) & \longrightarrow & H^{n+1}(L_1, \mu_m) & \longrightarrow & H^{n+1}(L_1, \mu_s) \\ \uparrow \text{res} & & \uparrow \text{res} & & \uparrow \text{res} \\ H^{n+1}(K, \mu_\ell) & \longrightarrow & H^{n+1}(K, \mu_m) & \longrightarrow & H^{n+1}(K, \mu_s) \end{array}$$

By Lemma 3.3, there is a field extension L of L_1 of degree at most ℓ so that L splits η . As a result, $\xi_L = \eta_L = 0$ and so L is a splitting field of ξ of degree

$$[L : K] = [L : L_1][L_1 : K] \leq \ell s = m.$$

The statement follows. \square

4. THE MAIN THEOREMS

In this section, we prove our two main theorems. First, we prove an analogue of Tate's common slot lemma over semiglobal fields. When considering coefficients in μ_ℓ for ℓ prime, we assume that $\text{cd}_\ell(k) = n$. For non-prime m , there is no notion of m -cohomological dimension and therefore we assume instead that $\text{cd}(k) = n$.

Let F be a semiglobal field with residue field k . We show in Theorem 4.3 that any finite set \mathcal{S} of r symbols in $H^{n+2}(F, \mu_m^{\otimes n+2})$ for m coprime to the characteristic of k is split by a common extension L of F of degree dividing m . We construct this extension explicitly in the proof of Theorem 4.3. First, we consider the case that $m = \ell$ is prime and we recall from Theorem 2.11 that

$$\text{III}_{\Omega_L}^{n+2} = \text{Ker} \left(H^{n+2}(L, \mu_\ell^{\otimes r}) \rightarrow \prod_{\omega \in \Omega_L} H^{n+2}(L_\omega, \mu_\ell^{\otimes r}) \right) = 0$$

for any integer r . By the above, it is sufficient to show that L splits any symbol $s \in \mathcal{S}$ locally. More precisely, let $\omega \in \Omega_L$ be a discrete valuation and denote its restriction to F by ν . We show that the restriction s_{F_ν} splits over L_ω . Finally, we extend the result to the case that m is any number coprime to the characteristic of k .

If $\mu_m \subset k$, then Theorem 4.3 implies that period and index agree in the top cohomology for semiglobal fields (see Proposition 4.5 for the precise statement). In the second part of this section, we remove the assumption that $\mu_m \subset k$ while retaining the equality for period and index in top cohomology. More precisely, in Theorem 4.8 we show that, for a semiglobal field F , any class ξ in $H^{n+2}(F, \mu_m)$ has index bounded above by m .

Recall the following setting for semiglobal fields. Let K be a complete discretely valued field and denote by \mathcal{O}_K its ring of valuations. Denote by C a smooth, projective curve over K , and $F = K(C)$ is the function field of C . In particular, F is a semiglobal field. Let k be the residue field of K . We discussed models and divisorial places in Section 2.3, but let us recall:

Definition 4.1 (Model). *A regular model of the curve C is a flat, projective, regular, integral \mathcal{O}_K -scheme \mathfrak{X} with $\mathfrak{X}_K = \mathfrak{X} \times_{\mathcal{O}_K} K = C$. We denote the reduced special fiber by \mathfrak{X}_k .*

$$\begin{array}{ccccc} C & \longrightarrow & \mathfrak{X} & \longleftarrow & \mathfrak{X}_k \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec}(K) & \longrightarrow & \text{Spec}(\mathcal{O}_K) & \longleftarrow & \text{Spec}(k) \end{array}$$

In the course of the proof, we will choose an appropriate model \mathfrak{X} so that the given cohomology class can only ramify along a divisor with normal crossings. Recall the notation Ω_F from Section 2.3 for the set of divisorial places on F , which is the set of valuations on F corresponding to codimension one points on all (regular, projective) models of C . To split the cohomology class, it will suffice to find an extension of F , which splits the cohomology class locally everywhere by Theorem 2.11.

We begin with the following. Let A be a complete regular local two-dimensional ring with field of fractions E and residue field k . For $m \geq 2$ and $1 \leq i \leq m$, we can write $a_i \in E^\times$ as

$$a_i = u_i \pi^{r_i} \delta^{s_i} \theta_i^\ell$$

for some units $u_i \in A$, integers $r_i, s_i \in \mathbb{Z}$ with $0 \leq r_i, s_i \leq \ell - 1$, and $\theta_i \in E$.

Lemma 4.2. *Let A be a complete regular local two-dimensional ring with field of fractions E and residue field k . Let (π, δ) be the maximal ideal of A , and take $a_i \in E^\times, 1 \leq i \leq m$. Then*

$$(a_1, \dots, a_m) = \xi_1 + (\xi_2, \pi) + (\xi_3, \delta) + (\xi_4, \pi, \delta),$$

where $\xi_1 \in H_{et}^n(A, \mu_\ell^{\otimes m})$, $\xi_2, \xi_3 \in H_{et}^{n-1}(A, \mu_\ell^{\otimes m-1})$, $\xi_4 \in H_{et}^{n-2}(A, \mu_\ell^{\otimes m-2})$ are unramified.

Proof. The proof follows along the same lines as the proof of Lemma 3.1. We remark that the assumption that $\mu_\ell \subset k$ in the proof of Lemma 3.1 is only required to prove the statement for a general cohomology class, which has to be written as a sum of symbols first. \square

We now prove an analogue of Tate's common slot lemma for number fields over semiglobal fields. We remark that the roots of unity μ_m may not be contained in k for this result.

Theorem 4.3. *Let F be a semiglobal field over a complete discretely valued field K , with residue field k . Let ℓ be a prime not equal to $\text{char}(k)$. Assume that $\text{cd}_\ell(k) = n$. Let*

$$\mathcal{S} = \{(a_{i,1}, a_{i,2}, \dots, a_{i,n+2}) : 1 \leq i \leq r\}$$

be a set of r symbols in $H^{n+2}(F, \mu_\ell^{\otimes n+2})$. Then there is a field extension L of F of degree ℓ which splits all symbols in \mathcal{S} .

Proof. We denote by $s_i = (a_{i,1}, \dots, a_{i,n+2})$ for $1 \leq i \leq r$ the symbols in \mathcal{S} . By Theorem 2.11, it suffices to prove the existence of a field extension L of F satisfying $[L : F] \leq \ell$ with the property that the restriction $(s_i)_{L_\omega}$ of s_i to L_ω is trivial for any divisorial place $\omega \in \Omega_L$. We will show the stronger statement that this is true for any discrete valuation ω of L , not just any divisorial valuation.

We now construct the desired splitting field L . Let \mathfrak{X} be any model of F so that the function field of \mathfrak{X} is F . We recall that \mathfrak{X}_k denotes the reduced special fiber of \mathfrak{X} . Abusing notation, we denote a representative of $a_{i,j}$ in F^\times by $a_{i,j}$ as well. As an element in the function field of \mathfrak{X} , $a_{i,j}$ is a rational function on \mathfrak{X} . Let $D_{i,j} = \text{div}(a_{i,j})$ be the corresponding divisor on \mathfrak{X} . After blowing up, we may assume that there are regular curves C_i so that

$$\mathfrak{X}_k \cup \left(\bigcup_{i,j} \text{supp}(\text{div}(a_{i,j})) \right) \subseteq \bigcup_i C_i$$

and the C_i have normal crossings. We denote $\mathcal{C} = \{C_i\}$. We call a point P on \mathfrak{X} a *nodal point of \mathcal{C}* if it lies on both C_i and C_j for some $i \neq j$. Choose a finite set of closed points $\mathcal{P} = \{P_j\}$ that includes all nodal points of \mathcal{C} and at least one point on the components of \mathfrak{X}_k . We note that even though the C_i may not be on the reduced special fiber \mathfrak{X}_k , all closed points P_j in \mathcal{P} lie on \mathfrak{X}_k .

Since \mathfrak{X} is projective, there is an affine open subscheme U of \mathfrak{X} that contains the finite set of closed points $\mathcal{P} = \{P_j\}$. We denote the affine coordinate ring of U by A . Since \mathfrak{X} is regular, A is regular as well. Let R be the semilocalization of A at \mathcal{P} . Because A is regular, we deduce that R is a semilocal regular domain and hence a unique factorization domain. Therefore each C_i defines a height one prime ideal in R , whose generator we denote by π_i . Finally, let $g = \prod_i \pi_i$ and set $L = F(\sqrt[r]{g})$. We aim to show that L is a splitting field for s_i for all $1 \leq i \leq r$. Any discrete valuation ω on L restricts to a discrete valuation ν on F . Let \mathcal{O}_ν denote the valuation ring of F at ν , F_ν the completion of F at ν and L_ω the completion of L at ω . We claim that L_ω is a splitting field for the restriction $(s_i)_{F_\nu}$ of s_i to F_ν for any $1 \leq i \leq r$. Throughout the remainder of the proof, we fix $1 \leq i \leq r$ and we denote $s = s_i$. We remark that by properness of the model \mathfrak{X} , any discrete valuation ν is centered on a closed point or on a point of codimension one. For any such P , we can define a regular local ring $\mathcal{O}_{\mathfrak{X},P}$. Let $m_{\mathfrak{X},P}$ be the maximal ideal of $\mathcal{O}_{\mathfrak{X},P}$, and denote by $\widehat{\mathcal{O}_{\mathfrak{X},P}}$ the completion of $\mathcal{O}_{\mathfrak{X},P}$ with respect to this maximal ideal.

- **Case 1: Suppose that ν is centered on a codimension one point P on \mathfrak{X} .** There are two possibilities for P , either $P \notin \mathcal{C}$ or $P \in \mathcal{C}$.
 - **Case 1a: Suppose that $P \neq C_i$ for all i .** By definition of the C_i 's and our hypothesis, a_{ij} is a unit in $\mathcal{O}_{\mathfrak{X},P}$ for all i and j . Thus s is unramified at the completion $\widehat{\mathcal{O}_{\mathfrak{X},P}}$ and so

$$s_{F_\nu} \in H_{\text{nr}}^{n+2}(F_\nu, \mu_\ell^{\otimes n+2}) \cong H_{\text{et}}^{n+2}(\widehat{\mathcal{O}_{\mathfrak{X},P}}, \mu_\ell^{\otimes n+2}) \cong H^{n+2}(k_\nu, \mu_\ell^{\otimes n+2}), \quad (4.1)$$

where the last isomorphism holds since $\widehat{\mathcal{O}_{\mathfrak{X},P}}$ is Henselian. We remark that, since P is a codimension one point, the residue field k_ν of F_ν

has transcendence degree 1 over k , and hence has the property that $\text{cd}_\ell(k_\nu) = n+1$. Using this and (4.1), we have that $s_{F_\nu} = 0$ and therefore its restriction s_{L_ω} also vanishes.

- **Case 1b: Suppose that $P = C_i$ for some i .** Since g is a parameter at C_i , $F_\nu(\sqrt[\ell]{g}) = L_\omega$, by Lemma 3.3, s_{F_ν} are split by the extension L_ω .
- **Case 2: Suppose that ν is centered on a codimension two point P on \mathfrak{X} .** Then $\mathcal{O}_{\mathfrak{X},P} \subset \mathcal{O}_\nu$ as a local homomorphism, which is an inclusion inducing an inclusion of the completions $\widehat{\mathcal{O}_{\mathfrak{X},P}} \subset \widehat{\mathcal{O}_\nu}$. As a result, the field of fractions of $\widehat{\mathcal{O}_{\mathfrak{X},P}}$ is contained in the field F_ν , which is a complete discretely valued field.

We distinguish three cases (2a) P is a nodal point of C_i and C_j for some $i \neq j$, (2b) P is a point on exactly one of the C_i , and (2c) P does not lie on any C_i . Denote by $k(P)$ the residue field of both $\mathcal{O}_{\mathfrak{X},P}$ and its completion.

- **Case 2a: Suppose that P is a nodal point on $\bigcup_i C_i$.** Since \mathfrak{X} has normal crossings, a nodal point will be at the intersection of two distinct curves C_i and C_j . Without loss of generality, assume that P is the intersection point of C_1 and C_2 . We remark that the maximal ideal of the local ring $\mathcal{O}_{\mathfrak{X},P}$ is generated by π_1 and π_2 and $\widehat{\mathcal{O}_{\mathfrak{X},P}}$ is a complete local ring of dimension 2. We also note that the residue field $k(P)$ is a finite field extension of k , thus $\text{cd}_\ell(k(P)) = \text{cd}_\ell(k) = n$.

By Lemma 4.2, we can decompose s_{F_ν} as

$$s_{F_\nu} = \xi_1 + (\xi_2, \pi_1) + (\xi_3, \pi_2) + (\xi_4, \pi_1, \pi_2),$$

for unramified cohomology classes $\xi_1 \in H_{et}^{n+2}(\widehat{\mathcal{O}_{\mathfrak{X},P}}, \mu_\ell^{\otimes n+2})$, $\xi_2, \xi_3 \in H_{et}^{n+1}(\widehat{\mathcal{O}_{\mathfrak{X},P}}, \mu_\ell^{\otimes n+1})$, and $\xi_4 \in H_{et}^n(\widehat{\mathcal{O}_{\mathfrak{X},P}}, \mu_\ell^{\otimes n})$. In conclusion, ξ_1, ξ_2, ξ_3 are unramified at $\mathcal{O}_{\mathfrak{X},P}$ and vanish over its completion as

$$H_{et}^{n+i}(\widehat{\mathcal{O}_{\mathfrak{X},P}}, \mu_\ell^{\otimes n+i}) \cong H^{n+i}(k(P), \mu_\ell^{\otimes n+i})$$

for $i = 1, 2$ and $\text{cd}_\ell k(P) = n$. We conclude that $s_{F_\nu} = (\xi_4, \pi_1, \pi_2)$. In $\mathcal{O}_{\mathfrak{X},P}$, we have that $g = u\pi_1\pi_2$ and so

$$\begin{aligned} s_{F_\nu} &= (\xi_4, \pi_1, \pi_2) \\ &= (\xi_4, -\pi_1, \pi_1\pi_2) \\ &= (\xi_4, -\pi_1, u^{-1}g) \\ &= (\xi_4, u, -\pi_1) + (\xi_4, -\pi_1, g) \\ &= (\xi_4, -\pi_1, g), \end{aligned}$$

where the last equality holds since (ξ_4, u) is unramified in

$$H^{n+1} \left(\widehat{\mathcal{O}_{\bar{x}, P}}, \mu_\ell^{\otimes n+1} \right) \cong H^{n+1} \left(k(P), \mu_\ell^{\otimes n+1} \right) = 0,$$

since $k(P)$ has cohomological dimension n . Using Remark 2.6, we see that $L_\omega = F_\nu(\sqrt[\ell]{g})$ splits s_{F_ν} .

- **Case 2b:** Suppose that P is a point on exactly one of the C_i . Without loss of generality, we may assume that P lies on C_1 . By Lemma 4.2, the class s_{F_ν} becomes

$$s_{F_\nu} = \xi_1 + (\xi_2, \pi_1)$$

for some $\xi_1 \in H_{et}^{n+2} \left(\widehat{\mathcal{O}_{\bar{x}, P}}, \mu_\ell^{\otimes n+2} \right)$ and $\xi_2 \in H_{et}^{n+1} \left(\widehat{\mathcal{O}_{\bar{x}, P}}, \mu_\ell^{\otimes n+1} \right)$. As in Case 2a, s is split by the degree ℓ extension given by attaching an ℓ -th root of $g = u\pi_1$, where π_1 is the local uniformizer at C_1 for ν .

- **Case 2c:** Suppose that P does not lie on C_i for any i . Then a_{ij} is a unit in $\mathcal{O}_{\bar{x}, P}$ and therefore s_{F_ν} is an unramified cohomology class in

$$H_{et}^{n+2} \left(\widehat{\mathcal{O}_{\bar{x}, P}}, \mu_\ell^{\otimes n+2} \right) \cong H^{n+2} \left(k(P), \mu_\ell^{\otimes n+2} \right).$$

The residue field $k(P)$ is a finite extension of k , implying that $\text{cd}_\ell(k(P)) = \text{cd}_\ell(k) = n$. Hence, the class $s_{\widehat{\mathcal{O}_{\bar{x}, P}}}$ vanishes, and so s_{F_ν} does as well. □

Corollary 4.4. *Let F be a semiglobal field over a complete discretely valued field K , with residue field k . Let $m \geq 2$ be coprime to $\text{char}(k)$. Assume that $\text{cd}(k) = n$. Suppose that $\mu_m \subset k$. Let*

$$\mathcal{S} = \{(a_{i,1}, a_{i,2}, \dots, a_{i,n+2}) : 1 \leq i \leq r\}$$

be a set of r symbols in $H^{n+2}(F, \mu_m^{\otimes n+2})$. Then there is a field extension L of F of degree ℓ which splits all symbols in \mathcal{S} .

Proof. We note that the assumption $\text{cd}(k) = n$ implies that $\text{cd}_\ell(k) \leq \text{cd}(k) = n$ for any prime ℓ dividing m . The proof follows from Theorem 4.3 and arguments analogous to the proof of Lemma 3.4. □

If k contains μ_m , the previous theorem implies that period and index agree in top cohomology. We later remove this assumption.

Proposition 4.5. *Let F be the function field of the smooth projective curve C over the complete discretely valued field K , with residue field k . Assume additionally that $\mu_m \subset k$, m is coprime to the characteristic of k , and $\text{cd}(k) = n$. Then any element ξ in $H^{n+2}(F, \mu_m)$ has index bounded above by m . In fact, there is a field extension L of F of degree m so that $\xi_L = 0$.*

Proof. Suppose $m = \ell$ is a prime number coprime to the characteristic of k and let $\xi \in H^{n+2}(F, \mu_\ell)$. By Voevodsky's result (see Section 2.2), we may write

$$\xi = \sum_{i=1}^r (a_{i,1}, \dots, a_{i,n+1})$$

for some $a_{i,j} \in F^\times$. We denote

$$\mathcal{S} = \{(a_{i,1}, a_{i,2}, \dots, a_{i,n+2}) : 1 \leq i \leq r\}.$$

By Theorem 4.3 there is some field extension L of F of degree ℓ which splits all symbols in \mathcal{S} . As a result, the sum ξ of the symbols in \mathcal{S} also splits over L . The general case m coprime to the characteristic of k with $\mu_m \subset k$ follows arguments by induction analogous to the proof of Lemma 3.4. \square

From now on, we no longer require that $\mu_m \subset k$. We first assume that $m = \ell$ is a prime number. Consider the field extension $F[\rho]$ obtained by attaching a primitive ℓ -th root of unity ρ . We denote by G the Galois group of $F[\rho]$ over F . To descend Theorem 4.7 to this setting, we first need to construct an appropriate model \mathfrak{X} for a fixed cohomology class $\xi \in H^{n+2}(F, \mu_\ell)$. Consider the restriction $\xi_{F[\rho]} \in H^{n+2}(F[\rho], \mu_\ell)$. Then $\xi_{F[\rho]}$ may be written as the sum of symbols

$$\xi_{F[\rho]} = \sum_{i=1}^r (a_{i,1}, \dots, a_{i,n+2})$$

for some $a_{i,j} \in F[\rho]$.

Lemma 4.6. *With the notation as above, there is a regular proper model \mathfrak{X} over \mathcal{O}_K of F such that the following conditions are satisfied:*

- (a) *The base change $\mathfrak{X}[\rho]$ of \mathfrak{X} to $\mathcal{O}_{K[\rho]}$ is a regular proper model of $F[\rho]$;*
- (b) *There are distinct irreducible curves C_1, \dots, C_r on $\mathfrak{X}[\rho]$ so that*

$$(\mathfrak{X}[\rho])_{k[\rho]} \cup \left(\bigcup_{i,j} \text{supp}(\text{div}(a_{i,j})) \right) \subseteq \bigcup_{\substack{1 \leq i \leq r \\ \sigma \in G}} C_i^\sigma, \quad (4.2)$$

where C_i^σ denotes the Galois conjugate of C_i ;

- (c) *For the projection $\pi : \mathfrak{X}[\rho] \rightarrow \mathfrak{X}$, we have that $\pi(C_i) = D_i$ for $1 \leq i \leq r$ are distinct irreducible curves on \mathfrak{X} ; and*
- (d) *$\sum_{i,\sigma} C_i^\sigma$ on $\mathfrak{X}[\rho]$ and $\sum_i D_i$ on \mathfrak{X} are divisors with normal crossings.*

Proof. Let \mathfrak{X} be a regular proper model of F over \mathcal{O}_K as discussed in Section 2.3 and denote by $\mathfrak{X}[\rho]$ its base change to $\mathcal{O}_{K[\rho]}$. We remark that the projection $\pi : \mathfrak{X}[\rho] \rightarrow \mathfrak{X}$ is a finite étale morphism. We may choose distinct irreducible curves D_1, \dots, D_r on \mathfrak{X} so that if C_1, \dots, C_r are distinct irreducible curves on $\mathfrak{X}[\rho]$ with $\pi(C_i) = D_i$, then

(b) is satisfied.

We remark that the model \mathfrak{X} does not necessarily satisfy (d). Let $b : \tilde{\mathfrak{X}} \rightarrow \mathfrak{X}$ be a blow up such that $b^*(\sum_{i=1}^r D_i)$ is a normal crossing divisor on $\tilde{\mathfrak{X}}$. We denote by $\tilde{\mathfrak{X}}[\rho]$ the base change, by $\tilde{\pi} : \tilde{\mathfrak{X}}[\rho] \rightarrow \tilde{\mathfrak{X}}$ the corresponding étale projection, and by $b[\rho] : \tilde{\mathfrak{X}}[\rho] \rightarrow \mathfrak{X}[\rho]$ the base change of b . Then the following diagram commutes

$$\begin{array}{ccc} \tilde{\mathfrak{X}}[\rho] & \xrightarrow{b[\rho]} & \mathfrak{X}[\rho] \\ \tilde{\pi} \downarrow & & \downarrow \pi \\ \tilde{\mathfrak{X}} & \xrightarrow{b} & \mathfrak{X} \end{array}$$

and therefore

$$\tilde{\pi}^* b^* \left(\sum_{i=1}^r D_i \right) = (b[\rho])^* \pi^* \left(\sum_{i=1}^r D_i \right) = (b[\rho])^* \left(\sum_{\substack{1 \leq i \leq r \\ \sigma \in G}} C_i^\sigma \right).$$

Since $b^*(\sum_{i=1}^r D_i)$ is a normal crossing divisor on $\tilde{\mathfrak{X}}$ and $\tilde{\pi} : \tilde{\mathfrak{X}}[\rho] \rightarrow \tilde{\mathfrak{X}}$ is finite étale, the pullback $(b[\rho])^* \left(\sum_{i,\sigma} C_i^\sigma \right)$ is a divisor with normal crossings on $\tilde{\mathfrak{X}}[\rho]$. This shows that the model $\tilde{\mathfrak{X}}$ satisfies the conditions (a)-(d) in the statement. \square

We are now ready to split the cohomology class $\xi \in H^{n+2}(F, \mu_\ell)$ using the model constructed in the previous lemma.

Theorem 4.7. *Let F be a semiglobal field over a complete discrete-valued field K with residue field k . Let ℓ be a prime number coprime to the characteristic of the residue field k , and suppose that $\text{cd}_\ell(k) = n$. Any element ξ in $H^{n+2}(F, \mu_\ell)$ is split in a field extension L of F of degree $\leq \ell$ so that $\text{ind}(\xi)$ divides ℓ .*

Proof. In view of Proposition 4.5, we may assume that F does not contain a primitive ℓ -th root of unity. Let ρ be a primitive ℓ -th root of unity in \bar{F} and remark that $[F[\rho] : F] = \ell - 1$. We write $\xi_{F[\rho]} = \sum_{1 \leq i \leq s} (a_{i,1}, \dots, a_{i,n+2})$, with $a_{i,j} \in F[\rho]$. Let \mathfrak{X} be a regular proper model chosen as in the previous Lemma, i.e. satisfying conditions (a)-(d). We denote $\mathcal{C} = \{C_1, \dots, C_r\}$ and $\mathcal{C}^\sigma = \{C_i^\sigma : 1 \leq i \leq r, \sigma \in G\}$, where G is the Galois group of $F[\rho]$ over F .

As in the proof of Proposition 4.5, we take $g \in F[\rho]$ so that

$$\text{div}_{\tilde{\mathfrak{X}}[\rho]}(g) = \sum_{i,\sigma} C_i^\sigma + E, \tag{4.3}$$

where E is a divisor on $\mathfrak{X}[\rho]$ whose support does not contain any C_i^σ and no component of $\text{supp}(E)$ passes through a nodal point of \mathcal{C}^σ . Let $N : F[\rho] \rightarrow F$ denote the norm map and we recall that $\pi : \mathfrak{X}[\rho] \rightarrow \mathfrak{X}$ denotes the projection. Let us also

denote $k(C)$ (resp. $k(D)$) the residue field at the codimension one point C on the 2-dimensional scheme $\mathfrak{X}[\rho]$ (resp. \mathfrak{X}). Then

$$\begin{aligned}
 \operatorname{div}_{\mathfrak{X}[\rho]}(N(g)) &= \pi^* \pi_* (\operatorname{div}_{\mathfrak{X}[\rho]}(g)) \\
 &\stackrel{(4.3)}{=} \pi^* \pi_* \left(\sum_{i,\sigma} C_i^\sigma + E \right) \\
 &= \pi^* \left(\sum_i f_i g_i D_i \right) + \pi^* \pi_* E \\
 &= (\ell - 1) \sum_{i,\sigma} C_i^\sigma + \pi^* \pi_* E,
 \end{aligned} \tag{4.4}$$

where f_i is the degree of the residue field extension $[k(C_i) : k(D_i)]$ and g_i is the number of conjugates C_i^σ of C_i . For the last equality, we remark that

$$f_i g_i = [F[\rho] : F] = (\ell - 1).$$

We now prove that no component of $\pi^* \pi_* E$ passes through a nodal point of \mathcal{C}^σ . Each component of $\pi^* \pi_* E$ is of the form E_r^τ , with E_r a component of $\operatorname{supp}(E)$ and $\tau \in G$. Suppose that E_r^τ passes through a nodal point $C_i^{\sigma_1} \cap C_j^{\sigma_2}$ of \mathcal{C}^σ . Then E_r passes through $C_i^{\sigma_1 \tau} \cap C_j^{\sigma_2 \tau}$ which is a nodal point of \mathcal{C}^σ contradicting that no component of E passes through a nodal point of \mathcal{C}^σ .

As in the proof of Theorem 4.3, let \mathcal{P} be a set of closed points on $\mathfrak{X}[\rho]$ that includes all nodal points of \mathcal{C} and at least one point on each component. We denote the set $\mathcal{P}^\sigma = \{P^\sigma : P \in \mathcal{P}, \sigma \in G\}$. Let U be an open subscheme of $\mathfrak{X}[\rho]$ that contains \mathcal{P}^σ and denote its affine coordinate ring by A . In the semilocalization R of A that corresponds to $U - \mathcal{P}^\sigma$ each C_i^σ defines a height one prime ideal whose generator we denote by π_i^σ . Without loss of generality, we may assume that $\sigma(\pi_i) = \pi_i^\sigma$. We denote $L = F \left(\sqrt[\ell]{N(g)} \right)$ and we claim that $L[\rho]$ splits $\xi_{F[\rho]}$. Let ω be a discrete valuation of $L[\rho]$ and denote its restriction to $F[\rho]$ by ν . If ν is centered on a codimension one point P on $\mathfrak{X}[\rho]$ so that $P \notin \mathcal{C}$, then the argument in the proof of Theorem 4.3 Case 1a applies to show that $\xi_{F[\rho]\nu}$ splits over $\xi_{L[\rho]\omega}$. If P is centered on C_i^σ for some i and some σ , then we remark that $N(g) = u (\pi_i^\sigma)^{f_i}$ for some unit $u \in \mathcal{O}_{\mathfrak{X}[\rho], C_i^\sigma}$. Since f_i is coprime to ℓ , the ℓ -torsion symbol (ξ_2, π_i) for $\xi_2 \in H^{n+1} \left(\widehat{\mathcal{O}_{\mathfrak{X}, C_i^\sigma}}, \mu_\ell \right)$ is split if and only if the symbol $(\xi_2, (\pi_i^\sigma)^{f_i}) = (\xi_2, \pi_i^\sigma)^{f_i}$ is split. Thus the argument in the proof of Theorem 4.3 Case 1b implies that $\xi_{F[\rho]\nu}$ splits over $L[\rho]\omega$. A similar reasoning shows the local splitting if ν is centered on a codimension-two point on $\mathfrak{X}[\rho]$. We conclude

that $L[\rho]$ splits $\xi_{F[\rho]}$. It remains to show that ξ is already split over L .

We consider the following commutative diagram

$$\begin{array}{ccc} H^{n+2}(L, \mu_\ell) & \xrightarrow{\text{res}} & H^{n+2}(L[\rho], \mu_\ell) \\ \text{res} \uparrow & & \uparrow \text{res} \\ H^{n+2}(F, \mu_\ell) & \xrightarrow{\text{res}} & H^{n+2}(F[\rho], \mu_\ell) \end{array}$$

The horizontal restriction maps are injective since $[F[\rho] : F] = [L[\rho] : L]\ell - 1$ is coprime to ℓ . Since ξ splits over $L[\rho]$, it also splits over the degree ℓ extension L of F by injectivity. \square

We note that we have therefore, in effect, proved the following theorem.

Theorem 4.8. *Let F be a semiglobal field over K a discrete valued field and denote k the residue field of the latter. Let m be any positive integer. Suppose that the characteristic of k is coprime to m and $\text{cd}(k) = n$. Then any element $\xi \in H^{n+2}(F, \mu_m)$ has index bounded above by m .*

Proof. Same argumentation as in Lemma 3.4, using Theorem 4.7 for the inductive assumption. \square

5. OPEN QUESTIONS

In this section, we discuss related results and open questions. We first propose a conjecture which extends our results on bounded index for the top cohomology for semiglobal fields. Secondly, we highlight the relationship between symbol length and the u -invariant for quadratic forms.

5.1. Questions on top cohomology. Let F be any field and fix a prime number ℓ that is coprime to the characteristic of F . We assume that $\text{cd}_\ell(F) = n$, so that $H^{n+r}(F, \mu_\ell) = 0$ for any $r > 0$. We call $H^n(F, \mu_\ell)$ the top cohomology of F .

We remark that in this generality, there is no uniform bound for indices of elements in top cohomology $H^n(F, \mu_\ell)$. More precisely, in [Mer91], Merkurjev produces examples of fields F with $\text{cd}(F) = 2$ which admit division algebras that are tensor products of n quaternion algebras for any given n . These algebras have index 2^n in $H^2(F, \mu_2)$. We are interested in uniform bounds for indices in top cohomology. As a first step in this program, we highlight the following conjecture.

Conjecture 5.1. *Let K be a field with $\text{cd}_\ell(K) = n$ and fix a prime ℓ coprime to the characteristic of K . Let F be a function field of a curve over K . If the indices in $H^n(K, \mu_\ell)$ are bounded, then the indices in $H^{n+1}(F, \mu_\ell)$ are bounded.*

We prove the conjecture if F is a semiglobal field. The conjecture also has a positive answer if K is a global field of characteristic different from ℓ [Sur20].

We recall that the diophantine dimension $\text{dd}(F)$ of a field F is the smallest integer $r \geq 0$ such that any n -form of degree $d \geq 1$ has a nontrivial zero in F , whenever $n > d^r$. If no such r exists, we say that $\text{dd}(F) = \infty$. The following conjecture was posed by Krashen in [Kra16, Conjecture 1, p. 997].

Conjecture 5.2. *Suppose that $\text{dd}(F) \leq d$. Then for elements $\xi \in H^d(F, \mu_\ell^{\otimes d})$, the index $\text{ind}(\xi)$ divides ℓ .*

Our main result may be viewed as an analogue of this conjecture with the assumption $\text{dd}(F) \leq d$ replaced by $\text{cd}(k) \leq d - 2$, which implies $\text{cd}(F) \leq d$, for a semiglobal field F .

5.2. Quadratic forms and u -invariant. It is well-known that classical invariants of quadratic forms take values in the Galois cohomology groups. In 1970, Milnor proposed higher invariants for quadratic forms which could determine the isomorphism class of a quadratic form up to hyperbolic planes.

Let $W(K)$ be the Witt ring of a field K and denote by $I(K)$ the fundamental ideal of $W(K)$ of even dimensional forms. The ideal $I^n(K)$ is generated by n -fold Pfister forms

$$\langle\langle a_1, \dots, a_n \rangle\rangle = \langle 1, -a_1 \rangle \otimes \langle 1, -a_2 \rangle \otimes \cdots \otimes \langle 1, -a_n \rangle.$$

For every $n > 0$, Milnor defined a map

$$I^n(K) \rightarrow H^n(K, \mu_2)$$

that takes the Pfister form $\langle\langle a_1, \dots, a_n \rangle\rangle$ to the symbol (a_1, \dots, a_n) . Milnor conjectured that these maps are well-defined and onto with kernel precisely $I^{n+1}(K)$. Milnor's conjecture was proved by Voevodsky in [Voe03].

Another invariant of fields associated with quadratic forms is the u -invariant which is defined as the maximum dimension of an anisotropic quadratic form over K . In the case that K is a complete discretely valued field with residue field k , Springer shows in [Spr55] that $u(K) = 2u(k)$. Here, we focus on the case of a semiglobal field F , which is the function field of a curve C over K . Suppose that $\text{char}(k) \neq 2$. If $u(l) \leq d$ for any finite extension l of the residue field k and $u(k(C)) \leq 2d$, where C is any curve over k , then $u(F) \leq 4d$ by [HHK09]. The case that K is a p -adic field was proved in [PS10]. On the other hand, if k is perfect with $\text{char}(k) = 2$, then Parimala and Suresh show in [PS14] that if F is the function field of a curve over a complete discretely valued field K of characteristic zero, then $u(F) \leq 8$. Further, they conjecture that with the exact same hypothesis, but that of k being perfect, $u(F) \leq 8[k : k^2]$. Equivalently, the question is whether every quadratic form in at

least nine variables over the function field of a curve over a local field has a nontrivial zero.

We now recall two main results in the area. The first of Krashen, [Kra16, 4.2], relates the bounds on indices of Galois cohomology classes for any field to bounds on the symbol length in Galois cohomology. The second result relates the bound on symbol length in μ_2 -cohomology to bounds on the u -invariant. Recall the notation from Section 2.2 on the symbol length of an element in $H^n(F, \mu_m)$ as $\text{symb}_n^m(F)$.

Theorem 5.3 ([Kra16]). *Let K be a field, ℓ a prime not equal to the characteristic of K and $n \geq 1$. Suppose that there exists an integer N such that for every finite extension L of K and for all $\beta \in H^d(L, \mu_\ell^{\otimes d})$, $1 \leq d \leq n$, $\text{ind}(\beta) \leq N$. Then, for any $\alpha \in H^n(K, \mu_\ell^{\otimes n})$, the symbol length of α is bounded in terms of $\text{ind}(\alpha)$, N and n .*

The following theorem is a consequence of a theorem of Orlov, Vishik, and Voevodsky [OVV07] on the Milnor conjecture (see also [PS15]).

Theorem 5.4 ([OVV07]). *Let K be a field of characteristic not equal to 2. Suppose that there exist integers $M \geq 1$ and N such that $H^M(K, \mu_2) = 0$ and $\text{symb}_d^2(K)(\alpha) \leq N$ for all $\alpha \in H^d(K, \mu_2)$, $1 \leq d < M$. Then the u -invariant is bounded by a function of M and N .*

If we could show that the index is bounded for elements in $H^i(F, \mu_2)$ for all $i \leq n + 2$, then using Theorems 5.3 and 5.4, it would follow that the u -invariant of F is bounded. In this paper, letting $\text{cd}(K) = n + 1$ for a discrete valued field K , we have shown that in the top cohomology (i.e degree $n + 2$) and in the context of a semiglobal fields F over K , the index is necessarily bounded. If we were to show that the index is bounded for all i -cohomology, $i \leq \text{cd}(K)$, then we would obtain a bound depending on this index for each $\alpha \in H^{n+1}(F, \mu_\ell^{\otimes n+1})$. Then applying Theorem 5.4, with $M = n + 2$, we would obtain a bound on the u -invariant as a function of three variables: the integer $n + 2$, some homogeneous bound on the indices for all finite extensions of F (in all degrees) and the index of the element in $H^{n+1}(F, \mu_\ell^{\otimes n+1})$.

Finally, we remark that the converse also holds: The finiteness of the u -invariant for μ_2 -cohomology implies the boundedness of the index in all degrees, as proven in [Sal15]. Further, if the u -invariant of F is bounded by 2^n where $\text{cd}_2(F) = n$, then by [Kra16, Proposition 5.2(ii)], the index of elements in $H^n(F, \mu_2)$ is at most 2.

REFERENCES

- [ABGV11] Asher Auel, Eric Brussel, Skip Garibaldi, and Uzi Vishne. Open problems on central simple algebras. *Transform. Groups*, 16(1):219–264, 2011.
- [BNH32] R. Brauer, E. Noether, and H. Hasse. Beweis eines Hauptsatzes in der Theorie der Algebren. *J. Reine Angew. Math.*, 167:399–404, 1932.
- [Cas86] J. W. S. Cassels. *Local Fields*. London Mathematical Society Student Texts. Cambridge University Press, 1986.

- [Gos22] Saurabh Gosavi. Generalized period-index problem with an application to quadratic forms, 2022. arXiv:1910.02473.
- [GS17] Philippe Gille and Tamás Szamuely. *Central simple algebras and Galois cohomology*, volume 165 of *Camb. Stud. Adv. Math.* Cambridge: Cambridge University Press, 2nd revised and updated edition edition, 2017.
- [HHK09] David Harbater, Julia Hartmann, and Daniel Krashen. Applications of patching to quadratic forms and central simple algebras. *Invent. Math.*, 178(2):231–263, 2009.
- [HHK14] David Harbater, Julia Hartmann, and Daniel Krashen. Local-global principles for Galois cohomology. *Comment. Math. Helv.*, 89(1):215–253, 2014.
- [HHK23] David Harbater, Julia Hartmann, and Daniel Krashen. Bounding cohomology classes over semiglobal fields, 2023. arXiv:2203.06770.
- [HKP21] David Harbater, Daniel Krashen, and Alena Pirutka. Local-global principles for curves over semi-global fields. *Bull. Lond. Math. Soc.*, 53(1):177–193, 2021.
- [Kra16] Daniel Krashen. Period and index, symbol lengths, and generic splittings in galois cohomology. *Bulletin of the London Mathematical Society*, page bdw060, 2016.
- [Lip75] Joseph Lipman. Introduction to resolution of singularities. In *Algebraic geometry (Proc. Sympos. Pure Math., Vol. 29, Humboldt State Univ., Arcata, Calif., 1974)*, pages 187–230. Amer. Math. Soc., Providence, R.I., 1975.
- [Lip78] Joseph Lipman. Desingularization of two-dimensional schemes. *Ann. of Math. (2)*, 107(1):151–207, 1978.
- [Mer91] A. S. Merkurjev. Simple algebras and quadratic forms. *Izv. Akad. Nauk SSSR Ser. Mat.*, 55(1):218–224, 1991.
- [NSW08] Jürgen Neukirch, Alexander Schmidt, and Kay Wingberg. *Cohomology of number fields*, volume 323 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2008.
- [OVV07] D. Orlov, A. Vishik, and V. Voevodsky. An exact sequence for $K_*^M/2$ with applications to quadratic forms. *Ann. of Math. (2)*, 165(1):1–13, 2007.
- [Pie82] Richard S. Pierce. *Associative algebras*, volume 9 of *Studies in the History of Modern Science*. Springer-Verlag, New York-Berlin, 1982.
- [PS98] R. Parimala and V. Suresh. Isotropy of quadratic forms over function fields of p -adic curves. *Inst. Hautes Études Sci. Publ. Math.*, (88):129–150 (1999), 1998.
- [PS10] Raman Parimala and V. Suresh. The u -invariant of the function fields of p -adic curves. *Ann. Math. (2)*, 172(2):1391–1405, 2010.
- [PS14] R. Parimala and V. Suresh. Period-index and u -invariant questions for function fields over complete discretely valued fields. *Invent. Math.*, 197(1):215–235, 2014.
- [PS15] R. Parimala and V. Suresh. On the u -invariant of function fields of curves over complete discretely valued fields. *Advances in Mathematics*, 280:729–742, 2015.
- [Sal97] David J. Saltman. Division algebras over p -adic curves. *J. Ramanujan Math. Soc.*, 12(1):25–47, 1997.
- [Sal98] David J. Saltman. Correction to: “Division algebras over p -adic curves” [J. Ramanujan Math. Soc. **12** (1997), no. 1, 25–47; MR1462850 (98d:16032)]. *J. Ramanujan Math. Soc.*, 13(2):125–129, 1998.
- [Sal15] David J. Saltman. Finite u invariant and bounds on cohomology symbol lengths. *Doc. Math.*, pages 577–590, 2015.
- [Ser73] Jean-Pierre Serre. *Cohomologie Galoisienne*, volume Vol. 5 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin-New York, 1973. Cours au Collège de France, Paris,

- 1962–1963, Avec des textes inédits de J. Tate et de Jean-Louis Verdier, Quatrième édition.
- [Ser97] Jean-Pierre Serre. *Galois Cohomology*. Springer Monographs in Mathematics. Springer-Verlag, New York-Berlin, 1997.
- [Spr55] T. A. Springer. Quadratic forms over fields with a discrete valuation. I. Equivalence classes of definite forms. *Indag. Math.*, 17:352–362, 1955. Nederl. Akad. Wetensch. Proc. Ser. A **58**.
- [Sur20] Venapally Suresh. Third Galois cohomology group of function fields of curves over number fields. *Algebra Number Theory*, 14(3):701–729, 2020.
- [Tat76] John Tate. Relations between K_2 and Galois cohomology. *Invent. Math.*, 36:257–274, 1976.
- [Voe03] Vladimir Voevodsky. Motivic cohomology with $\mathbf{Z}/2$ -coefficients. *Publ. Math. Inst. Hautes Études Sci.*, (98):59–104, 2003.

(DIJOLS) UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z4, CANADA
Email address: sarah.dijols@math.ubc.ca

(PARIMALA) EMORY UNIVERSITY, ATLANTA, GA 30322, U.S.A.
Email address: parimala.raman@emory.edu

(RAMDORAI) UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z4, CANADA
Email address: sujatha@math.ubc.ca

(URE) ILLINOIS STATE UNIVERSITY, NORMAL, ILLINOIS 61790, U.S.A.
Email address: cure@ilstu.edu